



ISSN: 2456-0057  
IJPNE 2016; 1(1): 46-52  
© 2016 IJPESH  
www.journalofsports.com  
Received: 11-02-2016  
Accepted: 12-03-2016

**Chandan Khanra**  
Mugberia Gangadhar  
Mahavidyalaya, Dept. of  
Nutrition, West Bengal, India.

## Food fortification in community health: Rural perspective of India

**Chandan Khanra**

### Abstract

Malnutrition in India, particularly among women, children and adolescents is an emergency needing immediate attention if the country has to have inclusive growth and development. Health portfolio tends to concentrate on infectious diseases (vaccination, oral rehydration, treatment of infections), and noncommunicable diseases like cancer and cardiovascular diseases. Important as these are, nutrition cannot be subsumed under these. Nutrition has to be the basis of judging national development. Without good nutrition, neither communicable nor non-communicable diseases can be controlled. Malnutrition is the worst form of noncommunicable disease and is an important risk factor for chronic diseases at a later date. Maternal malnutrition has multigenerational adverse effects on human health and development. Nutrition Security implies 'Physical, economic and social access to balanced diet, clean drinking water, safe environment, and health care'. Nutrition literacy and leadership at all levels is needed to understand and act. After 66 years of independence, India has among the highest incidence of under-nutrition in the world. Almost 50% of children under 5 are under weight (weight for age) and stunted (height for age). Over 30% of adults are also undernourished. Besides deficiency of calories and protein, deficiency of micronutrients (MN) (vitamins and minerals) is rampant. MN deficiency is referred to as the hidden hunger since often times it is not an obvious killer or crippler, but extracts heavy human and economic cost. Though anthropometric deficits are attributed to protein calorie malnutrition, MN deficiencies contribute significantly, because MNs are needed for utilization of proteins and calories and to fight infections from a young age.

**Keywords:** Food fortification, community health, rural perspective, Malnutrition

### Introduction

Food fortification is the process of adding vitamins and nutrients to food and drinks. The term *food fortification* is often used to describe one of two main processes. The first process, also referred to as food enrichment, is the adding back of nutrients to foods and drinks that have been removed during processing. The second process is the addition of nutrients to foods that are either not naturally occurring in the food or are added at a higher concentration than what is naturally occurring.

While the idea of fortifying foods has been around for centuries, in the modern era, the first widespread practice of food fortification began with the addition of iodine to salt in the early 20th century. During that time period, goiter, an enlargement of the thyroid gland, was a common condition in areas that did not have high concentrations of iodine in the soil. Public health officials discovered that the consumption of iodine could prevent goiter and found that the addition of iodine to salt was a simple yet effective solution to the problem.

### India - Fortification Status

#### Definition of the legislative status is

- **Mandatory:** Legislation that has the effect of mandating fortification of one or more types of wheat or maize flour or rice with at least iron or folic acid.
- **Planning:** There is written evidence that the government is acting to prepare, draft, and/or move legislation for mandatory fortification.
- **Voluntary:** Most countries allow voluntary fortification. We use this category if at least 50% of the industrially-milled wheat or maize flour or rice produced in the country is being fortified through voluntary efforts.

**Correspondence**  
**Chandan Khanra**  
Mugberia Gangadhar  
Mahavidyalaya, Dept. of  
Nutrition, West Bengal, India.

The fortification in India does not reach 50% of the state's total production, so it does not meet this definition of voluntary.

- **No fortification:** None of the above.

To keep up in today's busy world, people are becoming master multi-taskers, and when it comes to keeping up with their daily nutritional needs, they expect their foods to multi-task as well. According to the 2009 *International Food Information Council (IFIC) Functional Foods/Foods for Health Consumer Trending Survey*, the great majority of Americans believe that food provides benefits beyond basic nutrition and are interested in how certain foods or food components can improve or maintain their health. Today, many people can identify a specific food and/or food component and its associated health benefit. Historically, food fortification, such as iodized salt or vitamin D-fortified milk, served as a public health measure to address population-wide nutrient deficiencies. Now, there are also calcium- and vitamin D-fortified juices, breads fortified with omega-3 fatty acids, and vegetable oil spreads with plant sterols available for health-conscious consumers searching for foods with additional health benefits. These types of foods contain added nutrients and ingredients that may promote or support overall health and wellness in a variety of ways across many different body systems including heart, bone, digestive, eye, and brain health; weight management; and increased energy and immune health, among others. Foods not only must meet consumer needs and preferences but also address nutrition, regulatory, safety, and technical constraints. The European Food Safety Authority (EFSA), World Health Organization (WHO) and several countries, including the United States, United Kingdom, Australia, New Zealand, and Canada, follow similar guiding principles when it comes to their fortification policies.

### Fortifying our World

Fortification and enrichment are terms used to describe the addition of nutrients to foods but are two separate concepts. Enrichment refers to the *restoration* of nutrients lost during the handling, processing, or storage of foods, and levels are generally based on Food and Drug Administration (FDA) standards of identity. Fortification refers to the voluntary addition of nutrients at levels beyond those naturally occurring in the food.

Globally, the decision to fortify products is left up to individual food manufacturers. Voluntary fortification is a common practice in many countries. Additionally, 50 countries including the United States, Canada, and Australia require mandatory fortification of certain staple foods with specific nutrient(s) to improve public health, such as the fortification of enriched flour with folic acid to reduce the risk of neural tube birth defects, and restrict the fortification of foods with certain nutrients such as vitamin D.

### Types of foods can be fortified

The indiscriminate addition of nutrients to foods and the fortification of fresh produce; meat, poultry, or fish products; is not allowed by the U.S. government and fortification of unprocessed foods is prohibited in European countries. This is meant to help consumers understand the nutritional value of foods from each food group. Also fortification of some types of foods such as sugars and some snack foods (e.g., candies and carbonated beverages) is discouraged, so that consumers are not encouraged to choose fortified foods that are inconsistent with achieving dietary guidelines.

### How much is too much

All food ingredients must be safe at the level of use and for the intended use. In some cases the FDA has established maximum levels of certain ingredients while in other cases, the laws and regulations in the US allow for a manufacturer to make a determination of the safety of the ingredient for a specific intended use based on the views of experts qualified by scientific training to evaluate the safety of substances added to food. Chronic excessive intake of some nutrients above upper limits set by the Institute of Medicine may be harmful so fortification levels are set low to ensure a wide margin of safety. Baseline levels and population intakes are continuously monitored through the National Health and Nutrition Examination Survey, which is a national study that evaluates the nutrition status of Americans, and through other clinical, biochemical, and functional indicators.

### Consequences of micronutrient deficiencies

Apart from human suffering due to morbidity and mortality, malnutrition in general and MN deficiencies in particular have a high economic cost. 'Productivity losses due to poor nutrition are estimated to be more than 10% of lifetime earnings for individuals, and 2-3% of GDP to the nation. Cost of treating malnutrition is 27 times more than the investment required for its prevention'. According to a panel of Nobel laureates, of the top 10 priorities selected for advancing global welfare using methodologies based on the theory of welfare economics, in Copenhagen Consensus, 2008, were in the area of nutrition - micronutrient supplements, micronutrient fortification, bio fortification, de-worming and other nutrient programmes at school and community level<sup>8</sup>. These are also needed to achieve the millennium development goals.

### Iron Deficiency Anaemia (IDA)

Moderate and severe IDA adversely affects immunity (resistance to fight infections), cognitive and motor development, physical performance (and hence productivity) and reproductive health: (premature birth, low birth weight and perinatal mortality). It is estimated that anaemia is the direct cause of maternal deaths in 20% and contributory cause in another 20%. Apart from dietary deficiency, helminthic infections, inhibitors of iron absorption, in the diet and repeated pregnancies (in women) also contribute.

### Iodine Deficiency Disorder (IDD)

Goitre is the clinical manifestation of iodine deficiency disorder. The functional consequences are: permanent brain damage, (cretinism, - mental retardation, and deaf mutism), reproductive failure, and decreased child survival. Milder deficiency also adversely affects mental development.

### Vitamin A Deficiency

The earliest ocular manifestation of vitamin A deficiency (VAD) is night blindness, and Bitot spots on the white of the eye. Severe vitamin A deficiency leads to keratomalacia (ulceration and sloughing of the cornea) and total blindness. Though keratomalacia is no longer a public health problem, night blindness is prevalent particularly in pregnant mothers and subclinical deficiency (low Serum levels of Vitamin A), is still encountered. In addition to the ocular manifestations, vitamin A deficiency has been shown to cause growth retardation, decreased resistance to infections, and even death. Opinion regarding efficacy of vitamin A supplementation in reducing mortality even in populations with sub-clinical vitamin A deficiency is however divided.

### **B-Complex Deficiencies**

Though there is marked dietary, biochemical and clinical evidence of riboflavin (vitamin B2) deficiency (metabolically a very important vitamin), it has not received adequate attention because its deficiency is neither a killer nor a crippler. Impaired psychomotor performance in school children and adults and impaired reproduction in animals associated with riboflavin deficiency has been reported. There is evidence of dietary and biochemical folic acid deficiency in India. It can cause megaloblastic anaemia due to impaired red cell maturation. Folic acid deficiency has also been implicated in congenital malformation (neural tube defects), Folic acid supplementation in early pregnancy or even pre-pregnant state has been shown to prevent it. Folic acid deficiency leads to raised levels of serum homocysteine – an independent risk factor for cardiovascular disease (CVD). Fragmentary evidence suggests that Indians do tend to have high levels of homocysteine which responds to treatment with folic acid<sup>16</sup>. Till recently, vitamin B12 deficiency was not considered to be a problem in India since its daily requirement is only 1 microgram. However, reports of vitamin B12 deficiency in developing countries like India and its link with homocysteinaemia, besides megaloblastic anaemia, have started appearing<sup>17</sup>. Both folic acid and B12, besides vitamin B6 and B2 are required for homocysteine metabolism.

In view of the rising incidence of CVD in India, B- complex vitamin deficiency needs to be taken more seriously and its link with homocysteinaemia and CVD needs to be investigated. Research is also needed to examine the role and dosage of folic acid for prevention of neural tube defects—which are not uncommon in India. A balance of folic acid with vitamin B12 has to be ensured.

### **Vitamin D Deficiency**

Main function of vitamin D is in bone calcification by facilitating calcium absorption and maintaining blood calcium levels. Since generation of vitamin D in the skin from its precursor 7-dehydrocholesterol is through exposure of skin to sunlight, adequacy of vitamin D in a tropical country like India was assumed. However, recent studies suggest existence of vitamin D deficiency in all age groups in India. As mentioned earlier, osteoporosis associated with calcium and vitamin D deficiency is common in post-menopausal women. Low levels of vitamin D are also associated with chronic diseases like certain malignancies, and chronic inflammatory and autoimmune diseases like type 1 diabetes, and impaired resistance to infections<sup>5, 9, 10</sup>. However, according to a recent review “Health—benefits often reported in the media—were from studies that provided often mixed and inconclusive results and could not be considered reliable”. Research is needed to determine the dietary requirement of vitamin D to ensure adequate vitamin D status.

### **Vitamin C Deficiency**

Vitamin C is a powerful antioxidant. Dietary vitamin C deficiency does exist, but severe clinical manifestation (scurvy) has become rare. Vitamin C is an iron absorption promoter and hence its deficiency can contribute to IDA. Antioxidants delay degenerative diseases.

### **Zinc Deficiency**

Zinc is essential for growth and development. Zinc supplementation has been reported to help linear growth, reduce severity and duration, of diarrhoeas, and respiratory infections and reduce child mortality. The magnitude and

consequences of zinc deficiency in India need to be determined. For the purpose suitable indicators have to be developed to assess zinc deficiency. Strategies for Increasing Access to Micronutrients — Current Response and Research Opportunities Basically there are four types of approaches to augment MN intake. 1) Pharmaceutical supplements, 2) Food fortification 3) Biofortification and 4) Food-food fortification (dietary diversification).

### **Micronutrient Supplementation**

#### **National Nutritional Anaemia Control Programme (NACP)**

In this programme supplements containing 100 mg of elemental iron + 500 µg folic acid are given to pregnant women for 100 days during pregnancy; 20 mg elemental iron and 100 µg folic acid are given daily to preschool children for 100 days in the year.<sup>9,10</sup>. Recently adolescent girls have also been included as part of the life cycle approach with same dose as pregnant women, and weekly once administration throughout the year. Unfortunately despite scientific basis for the programme, iron folic acid supplementation has failed to have an impact on the incidence or severity of anaemia, allegedly due to 1) lack of awareness regarding its importance and consequently poor compliance and 2) poor outreach in a vast country like India. NFHS 3 survey shows less than 20% full compliance in pregnant and lactating women.<sup>4</sup> 3) uniformly giving one tablet of IFA (which is meant for preventing anaemia in non-anaemic women) regardless of the severity of anaemia. Screening of all pregnant women for anaemia and treatment of anaemic women besides the prophylactic treatment with iron, folic acid has been recommended to reduce the prevalence and severity of anaemia in pregnancy.

#### **Suggestions for Research to Improve NACP**

Public-private-NGO partnership may have a role in improving the outreach. Absorption of non-haeme Iron from the diet is only 3-5%. The challenge is to translate filling of iron stores in to improvement in haemoglobin.

Research is needed to find out socio-cultural, behavioural factors and administrative bottle neck to improve the efficiency of NACP.

All pregnant women should get Hb estimation done using reliable method and anaemic women treated with appropriate route and dose of iron, folic acid.

Children coming to hospital for any illness and undernourished children should be screened and those found anaemic should be appropriately treated.

Prevalence of iron, folic acid and B12 deficiency in non anaemic and anaemic pregnant women in different regions of the country should be assessed to find out if the current practice of prescribing iron and folic acid without B12 is appropriate.

Efficacy of present regimen of giving uniformly one tablet of iron –folic acid needs re-evaluation and replaced with treatment after screening.

Since nutrients, besides iron and folic acid, are also involved in haemoglobin synthesis/formation of red blood cells/absorption of iron, inclusion of MN like vitamins B12, C, B2, and zinc may improve the efficacy of the oral supplements.

Clinical research in a hospital setting with appropriately worked out dose and schedules for multi-vitamin supplements Vs IFA supplements should be conducted.

### **Massive Dose Vitamin A Supplementation to Prevent Nutritional Blindness**

In this programme children between 6-60 months are given 200,000 IU of vitamin A, every six months as prophylactic dose. The rationale is: vitamin A being fat soluble, is stored in the liver and a massive dose would ensure adequate storage to last for at least 6 months. This programme also suffers from the infirmity of the other programmes viz., poor outreach, inadequate and irregular supplies. As mentioned earlier the severity of vitamin A deficiency has reduced despite inefficient operation of this programme and several eminent nutrition scientists have raised doubts about its continuation. However, as of now, this programme should continue in areas where vitamin A deficiency is a public health problem (incidence of Bitot spots more than 0.5%).

According to the Annual Report of Micronutrients Initiatives India, an International Non-Government Organization, out of 32 Million US Dollars available in the Annual Budget 2009-2010, more than 20 Million US Dollars were spent on Vitamin A Procurement and Interventions. A meager sum of 2.5 Million US Dollars was spent on Iron interventions (Umesh Kapil, personal communication). Children with VAD should be identified and should be administered VA. The present Universal VAS approach needs to be reviewed.

### **Food Fortification**

#### **Iodised Salt**

Food fortification is a powerful method of reaching out a deficient nutrient to populations, provided the vehicle used for fortification is consumed by the poorest of the poor. In India salt is the only such vehicle and it has been effectively used for reaching out iodine. This is one successful programme in the country, but its efficiency has to be improved in terms of stability of iodine in the salt, pricing, and outreach. Private companies tend to seek easy urban markets even if the problem is more acute elsewhere.

#### **Iron Fortified Iodised Salt (IFIS, also called Double Fortified Salt-DFS)**

This technology was developed by the National Institute of Nutrition, Hyderabad to address the dual problem of iron and iodine deficiency. Government order for its production has been released. Its efficacy has been tested in small scale studies.

Systematic programme for scaling up is needed to examine the effectiveness of DFS in preventing iron and iodine deficiency and replacing iodised salt with IFIS. Evaluation of DFS should be done by an independent agency rather than ICMR.

#### **Iron Fortified Wheat Flour (Atta) and Rice**

Since staple grains are consumed in substantial quantity, their fortification makes sense. In some countries wheat flour is fortified with iron and other micronutrients. Doubts have been raised about bio-availability of iron from wheat 'atta' because of high phytate (inhibitor of absorption) content. The inhibitory effect of phytate may be bypassed by some potential compounds like Na-Fe-EDTA and or enzyme phytase. The higher cost of this salt may be off-set by better bioavailability and hence lesser dose of fortification.

For more than a half the population in India, rice is the staple. Fortification of rice has been tried by mixing fortified extruded grains from rice flour with rice (Ultra rice). More research is needed to make this technology cost-effective and acceptable. Programmatic studies are needed to examine the effectiveness of fortifying cereals with iron. Fortification of Cereal Products

with Folic Acid In many countries, cereal products are fortified with folic acid to reduce the incidence of neural tube defects. Folic acid fortification, perhaps along with vitamin B12 may also reduce serum homocysteine levels.

In view of the rising incidence of CVD in India, this strategy needs to be researched.

#### **Fortification of oil with Vitamins A and D**

Gujarat has taken the initiative. Impact studies are needed. Fortified oil should be packed in dark bottles to cut off UV radiation, and prevent oxidation of vitamin A

Vehicles like sugar or soya sauce used in some countries are not suitable for India.

#### **Fortification of Food for Supplementary Feeding**

India has large feeding programmes like supplementary feeding of preschool children through ICDS and school children through Mid-Day Meal. Opinion of nutritionists is divided about fortifying these foods with MNs – sprinklers (micronutrient powder- MNP), spreads.

Well planned field research to examine efficacy, feasibility and cost of fortification vs. enrichment with MN rich foods, of meals for supplementary feeding need to be undertaken. Impact of Golden rice rich in pro vitamin A, ultra rice rich in iron and other MN and red palm oil in feeding programmes needs to be researched for feasibility, acceptability and cost effectiveness.

Food fortification is suitable only for prevention of MN deficiencies and not for treating severe forms of the disease. Food fortification programme for country should consider the Recommended Dietary Allowances (RDA) of different nutrients for Indians, the types of foods to be fortified, nutrients to be added and the percentage of the RDA to be supplied through fortification etc. to avoid excess intakes by the rich and deficient intakes by the poor. Fortified foods should be by the poorest of the poor, and reach the unreachable in a vast country like India.

#### **Bio-fortification**

Enriching germplasm with MNs through conventional breeding methods, molecular breeding and genetic engineering is a promising method for increasing dietary access to MNs. Bio-fortification is a sustainable intervention being a seed-based technology. There is no recurring cost, once the varieties are developed and adopted. It can benefit the farmer as well as the consumer if the cost of seed is kept low and not exploited by seed companies. Bio-fortified plants grow better. Potential nutritional impact of iron biofortification in India seems to be encouraging. "The Harvest Plus: bio-fortification challenge programme is an interdisciplinary, global alliance of research and implementing institutions. India is part of this. It includes:  $\beta$  carotene (pro-vitamin A)- rich sweet potato, and cassava, zinc and iron- rich rice, wheat, maize, pearl millet, and beans. DBT network project on biofortification of rice, wheat and maize is currently being implemented by ICAR Institutions and state agriculture universities and National Institute of Nutrition. Golden Rice rich in pro-vitamin A; high- iron rice (high ferritin gene from mangrove); are examples of transgenic technologies."

For conventional breeding or molecular breeding to improve the MN content of foods, sufficient within species diversity with appropriate gene pool would be necessary. Also this approach is slow. GM route with genes from other plant species, preferably edible varieties, is faster and with suitable safe guards regarding safety to health, protection of

biodiversity, and cost; is a useful option for a country like India. While the conventional strategies of food fortification and plant breeding to improve nutritional quality should be pursued, the GM technology despite some concerns needs to be vigorously investigated. This technology has received support from many science academies of the world including Indian academies.

#### **Food–Food Fortification—Dietary Diversification**

The problem of MN deficiency can to a great extent be addressed by encouraging dietary diversification and household access to MN-rich foods.

Overall emphasis of agriculture, horticulture and livestock breeding has been to ensure calories, income and export. MNs are not on the antenna of agriculture planning. Decentralised planning to ensure district, village and household level adequacy of MN- rich foods is required. India being a vast country, long distance transport of perishable foods like vegetables and animal products may not be feasible. Promotion of homestead gardens even in urban areas, with emphasis on MN- dense varieties, back- yard poultry, dairy, fish ponds etc is needed.

Operation research to study the impact of such a strategy needs high priority. Farm-based approaches are low-cost, and sustainable if the community is empowered.

While it may be difficult to meet the requirement of some nutrients like iron and riboflavin through low-cost diets, there is ample supply of  $\beta$  carotene (pro-vitamin A) in nature in green leafy vegetables and yellow orange fruits and vegetables. This wealth needs to be exploited through education and advocacy. Low bio-availability of beta carotene is an issue. Food habits may be hard to change.

Research is needed to establish the factor for conversion of  $\beta$  carotene to vitamin A and improve its bioavailability. Red palm oil is an important source of  $\beta$  carotene and can be used in supplementary feeding programmes if supply and acceptability can be assured.

Absorption and utilisation of micronutrients is influenced by competition between nutrients, (particularly minerals and trace elements), and other biotic (microorganisms, helminthic) and a-biotic (chemical agents like phytates, tannins, xenobiotics etc.).

These influences have to be researched to improve utilisation of dietary MNs. Treatment with phytases and tannases needs to be considered.

Bioavailability studies to examine the interaction within nutrients, and between nutrients and promoters, inhibitors and xenobiotics (biotic and abiotic), have to be planned.

#### **Enhanced Production to Consumption of Millets and Pulses**

MN rich coarse grains and millets are being forgotten. They are referred to as orphan grains, but with the threat of global warming, they may well be the grains of the future. There is 2-3 fold gap between optimal productivity of millets and farm-level productivity. Apart from being high in proteins, pulses are also rich in MN. Their production and consumption is on the decline.

Productivity of these grains has to be increased through research and robust extension effort. Agriculture research needs to give high priority to millets and pulses – a production to consumption strategy. Some of the World Bank funded National Agriculture Innovation Projects (NAIP) are attempting to do that.

#### **Plant foods as specialty foods for protection against chronic diseases**

This is a vast field for research since plant foods are rich in health giving phytochemicals-nutraceuticals. Some nutrients like vitamin C, E, Se also have antioxidant properties.

#### **Economic Logic**

World Bank has compared the benefits of MNs (iron) supplementation vs. Food fortification in terms of 1) cost per life saved, 2) productivity gained- a measure of efficient use of resources and defined as least cost discounted method of reducing clinical deficiency in the population and 3) social benefit cost- DALY or healthy life years saved. For saving life at least cost, targeted iron supplementation to pregnant women is more effective than iron fortification of flour or other staple. However, for enhanced productivity delivered by a programme and social benefit cost, iron fortification is most cost effective. Similar analysis is needed in the Indian context. Farm-based interventions are believed to have high benefit-cost ratio in terms of economic cost. They also help the farmer and empower the community.

#### **Food Analysis**

India is rich in biodiversity with lots of local foods with potential benefits in terms of MN as well as health-promoting phytochemicals. Food analysis labs equipped to do co-ordinated MN analysis with proper quality control should be set up in agriculture universities and home science colleges which have the expertise and country-wide research programmes to analyse local foods and identify MN rich foods should be started so that their production and consumption can be enhanced.

#### **Food Storage and Processing**

Almost 30% of farm food perishes due to lack of storage facilities and value addition. Development of suitable storage structures not only for grains but also for vegetables, fruits and animal products and promoting those which are already available is urgently needed. There are indigenous time-tested traditional technologies for storage. Those have to be re-discovered and promoted.

Value addition should not take away food from the poor. Both, high-end and low-cost, processed foods need to be developed with public private partnership, with the triple objectives of reducing wastage, generating employment and enhancing nutrition value and security.

#### **Bioavailability and Stability of Micronutrients in Processed Foods**

While minerals are reasonably stable during processing, vitamins, particularly vitamins A and C are sensitive to light and heat. Microencapsulation is done to increase the stability, but that increases the cost. Low-cost technologies to increase the bioavailability and stability of MNs in processed foods are needed to make processed foods accessible to all sections of the society.

#### **Nutrition Security and Climate Change**

Rise in temperature due to climate change is associated with increase in atmospheric carbon dioxide. While the latter helps to increase plant growth if there is enough supply of other nutrients in the soil, in tropical climate, further temperature rise has an adverse effect. It is suspected that apart from plant growth, climate change may also affect the nutrient composition (protein and minerals) in foods. No information is

available on vitamin content of foods and climate change. Impact of climate change on productivity and nutrient composition of major food items in India needs to be researched.

### Monitoring, Surveillance and Management Information System for Early Detection

India needs a robust monitoring and surveillance programme in nutrition for early identification of the problems and capacity building to tackle them in a decentralized way. For every frank case of deficiency, there are dozens of others who suffer from subclinical deficiencies which contribute substantially to human suffering, medical expenses and economic cost. Sensitive biochemical and functional tests to detect such deficiencies, using new knowledge of molecular and cellular biology and state of the art instruments need to be developed

### Nutrition and Health Education

Behavioural Change Communication (BCC) programs to targeted audience can improve both nutrition and health status. Nutrition education for professionals (agriculture, medical, social scientists and others), needs to be strengthened. Innovative strategies need to be developed and tested not only to improve knowledge and attitudes but practices as well. Behavioural modification modules are needed. Nutrition education policy is nonexistent. Awareness among school children, teachers, consumers and women has to be enhanced. Women empowerment is essential to improve diets and health. Nutrition component of medical and agriculture curricula needs to be strengthened.

### Conclusion

The current mindset of looking at food security only in terms of energy security has to change. Pumping cereals alone to quench hunger will not ensure nutrition and health. The goal should be to ensure a balanced diet adequate in macro- and micronutrients. Laboratory, clinical, and community (operations) -based research is needed to ensure MN security. An optimum mix of food-food fortification, (dietary diversification), bio fortification, and early detection and effective treatment of clinical deficiencies needs to be worked out. Extension methodology has to be robust. Media support for creating awareness and compliance is important. The most important low-cost intervention is following WHO guidelines for breast feeding- early initiation, exclusive breast feeding for 6 months and complimentary feeding with continuation of breast feeding up to 1 year. This requires media blitz to educate the community.

### References

1. UNICEF State of World's Children, Report, 2010, special edition
2. National Nutrition Monitoring Bureau (NNMB), Prevalence of micronutrient deficiencies; Technical report No.22, National Institute of Nutrition, ICMR, 2003
3. National Nutrition Monitoring Bureau (NNMB), Diet and nutrition status of populations and prevalence of hypertension among adults in rural areas;. Technical Report No 24. National Institute of Nutrition, ICMR, 2006
4. National Family Health Survey (NFHS-3), International Institute of population Sciences, Mumbai, India, 2005-2006; Jan.2011, 96 (1).
5. Ross C, Manson JE, Abrams SA *et al.* The 2011 Report on dietary reference intakes for calcium and vitamin D

6. from the Institute of Medicine: What clinicians need to know; J Clin Endocrinol Metab,96:53-58, 2011
6. Research and development in millets. Present status and future strategies. National seminar on millets-2010; Directorate of sorghum research, Hyderabad, November, 2010.
7. Coalition for sustainable nutrition security for India, Leadership agenda, 2010; <http://www.nutritioncoalition.in/publication.asp>
8. Copenhagen Consensus,2008.<http://www.copenhagenconsensus.com/Home.aspx>.
9. Textbook of human nutrition, 3rded. (2010). Editors: Bamji MS, Krishnaswamy, K and Brahmam GNV
10. Nutrient requirement and recommended dietary allowances for Indians. A report of the expert group of the Indian Council of Medical Research, 2010; <http://www.icmr.nic.in/pricepubl/content/11.htm>
11. Latham M. The great vitamin A fiasco; World Nutr; 1:12-45, 2010,
12. Gopalan C. Massive dose vitamin A prophylaxis should now be scrapped;. World Nutr.,1:79-85. 2010
13. Kapil U and Sachdev HPS. Universal Vitamin A Supplementation Programme in India: The need for a re-look; The National Medical Journal of India.23:193-195, 2010
14. Bamji MS and Lakshmi AV, Less recognised micronutrient deficiencies in India; NFI Bulletin, 19:5-8, 1998
15. Lakshmi AV and Bamji MS. Aetiology and health implications of hyperhomocysteinaemia; PINSA B64:235-248, 1998
16. Joshi S. Demystifying relevance of homocysteine hypothesis in native Asian Indian population. Editorial. J. Assoc. physicians, India; 54:763-764, 2006
17. Stabler S. and Allen RH. Vitamin B12 deficiency as a worldwide problem; Annual Review of Nutrition, 24:299-326,2004
18. Ranganathan S and Sesikaran B. Development of the double fortified salt from the National Institute of Nutrition; Comprehensive Rev Food Sci, Food Safety.7:390-396,2008.
19. Transforming Development through Science, Technology and Innovation. Ultra Rice Challenge; <http://www.usaid.gov/scitech/ur.html>. (Accessed January 3, 2011).
20. Stein A, Meenakshi J, Qaim, JV, et.al. Potential impacts of iron bio-fortification in India; Social Science and Medicine, 66:1797-1808, 2008
21. Harvest plus in Asia. Breeding crops for better nutrition. [harvestplus@cgiar.org](mailto:harvestplus@cgiar.org). [www.harvestplus.org](http://www.harvestplus.org). Accessed Jan 3, 2011
22. "Nutrition Security for India- issues and the way forward" INSA Position paper, [www.insa.ac.in](http://www.insa.ac.in)
23. Hunt JM, Forging effective strategies to combat iron deficiency; J. Nutr. 132: 794S-801S, 2002
24. Manoj-Kumar and Patra AK. Nutritional security: a missing link in climate change debates; Current Science, 99:287-289, 2010

### Reviewers

1. Dr C Gopalan, President, Nutrition Foundation of India, New Delhi
2. Professor MS Swaminathan, Chairman – MS Swaminathan Research Foundation, Chennai

3. Professor PN Tandon, President, National Brain Research Centre, New Delhi
4. Professor MS Valiathan, National Research Professor, Manipal University, Manipal
5. Professor G. Padmanabhan, Platinum Jubilee Chair, and Hon. Professor, IISc, Bangalore
6. Dr T Ramasami, Secretary, Department of Science and Technology, New Delhi
7. Professor RP Sharma, INSA Sr. Scientist, RCPB, ICAR, New Delhi
8. Professor Akhilesh Tyagi, Director, NIPGR, New Delhi
9. Professor Jacob John, Former Professor, CMC, Vellore
10. Dr Kamala Krishnaswamy, Former Director, National Institute of Nutrition, Hyderabad
11. Dr Vinodini Reddy, Former Director, National Institute of Nutrition, Hyderabad
12. Dr Prema Ramachandaran, Director, Nutrition Foundation of India, New Delhi
13. Professor V Prakash, Director, CFTRI, Mysore
14. Dr B Sivakumar, Former Director, National Institute of Nutrition, Hyderabad
15. Dr B Sesikaran, Director, National Institute of Nutrition, Hyderabad
16. Dr GNV Brahmam, Scientist F, National Institute of Nutrition, Hyderabad
17. Dr M Raghunath, Scientist E, National Institute of Nutrition, Hyderabad
18. Dr K Madhavan Nair, Scientist E, National Institute of Nutrition, Hyderabad
19. Dr S Ranganathan, Former Head, Isotope Division, National Institute of Nutrition, Hyderabad
20. Professor Umesh Kapil, Department of Gastroenterology and Human Nutrition Unit, AIIMS, New Delhi
21. Dr Tara Gopaldas, Former Professor, Dept. of Food and Nutrition, MS University, Baroda Professor Subhadra Seshadri, Former professor and Head, Dept. of Food and Nutrition, MS University, Baroda
22. Professor Venu Seth, Lady Irvin College, New Delhi
23. Professor Anura Kurpad, Dean St. Johns Research Centre, Bangalore
24. Dr Klaus Kraemer, Editor, Sight and Life, Switzerland