



ISSN: 2456-0057

IJPNE 2019; 4(1): 236-238

© 2019 IJPNE

www.journalofsports.com

Received: 17-11-2018

Accepted: 20-12-2018

Aabid UR Rehman

PhD, Scholar, Dept. of Physical Education and, Annamalai University, Annamalaiagar, Tamil Nadu, India

Dr. Ponson

Assistant Professor, Dept. of Physical Education, Annamalai University, Annamalaiagar, Tamil Nadu, India

Impact of submaximal and maximal aerobic training on selected pulmonary variable among under 14 years cricketer

Aabid UR Rehman and Dr. Ponson

Abstract

The purpose of the study was to find out the impact of submaximal and maximal aerobic training on selected pulmonary variable among under 14 years cricketer. Forty five under 14 years cricketer were selected randomly as subjects from YMCA, Cricket Academy at Chennai city, Tamil Nadu, India. The subjects were divided in to three groups of fifteen each named as submaximal aerobic training group, maximal aerobic training group and control group. The training program is scheduled at 6.30 am to 7.30 am on alternate days. The VO_2 max was selected as criterion variables and tested by Queens's college step test. The subjects were tested prior to and after the twelve weeks of experimentation. The obtained data from the experimental and control group were statistically analyzed with analysis of covariance (ANCOVA). Scheffe's post hoc test applied to examine the paired mean difference between groups if, the obtained 'F' value found significant on the selected criterion variable. The level of confidence was fixed at 0.05. The result shows that the experimental groups had achieved significant improvement on VO_2 max when compared to control group.

Keywords: Sub-maximal aerobic, maximal aerobic, pulmonary and VO_2 max

Introduction

Modern-day cricket has experienced a shift towards limited over games, where the emphasis is on scoring runs at a rapid rate. Although the use of protective equipment in cricket is mandatory, players perceive that leg guards, in particular, can restrict their motion Physical demands of cricket presumably vary by both game format and performance level. Cricket is the most popular sport in India; it is played by many people in open spaces throughout the country though it is not the nation's official national sport

"Maximal oxygen uptake (VO_2 max) is widely accepted as the single best measure of cardiovascular fitness and maximal aerobic power. Absolute values of VO_2 max are typically higher in men than in women. The average untrained healthy male will have a VO_2 max of approximately 35–40 mL/(kg· min). These score can improve with training and decrease with age, though the degree of trainability also varies very widely: conditioning may double VO_2 max in some individuals. In sports where endurance is an important component in performance, such as cycling, rowing, cross-country skiing, swimming and running, world-class athletes typically have high VO_2 max. (Heyward V, 1998) [12].

The term fitness is an important aspect to be developed in the minds of all the people irrespective of age and sex. Much attention has to be focused on youth physical fitness. A sound and well organized physical education program in the schools and colleges will be right solution for these problems. (Bucher, 2002).

Methodology

To achieve the purpose of the study, forty five under 14 years cricketer were selected randomly as subjects from YMCA, Cricket Academy at Chennai city, Tamil Nadu, India. The subjects were divided in to three groups of fifteen each named as submaximal aerobic training group, maximal aerobic training group and control group. The training program is scheduled at 6.30 am to 7.30 am on alternate days. The VO_2 max was selected as criterion variables and tested by Queens's college step test.

Correspondence

Aabid UR Rehman

PhD, Scholar, Dept. of Physical Education and, Annamalai University, Annamalaiagar, Tamil Nadu, India

The subjects were tested prior to and after the twelve weeks of experimentation. The obtained data from the experimental and control group were statistically analyzed with analysis of covariance (ANCOVA). Scheffe's post hoc test applied to examine the paired mean difference between groups if, the

obtained 'f' value found significant on the selected criterion variable. The level of confidence was fixed at 0.05.

Results and discussions

Table 1: Analysis of covariance on vo2 max of experimental and control group

Mean	Submaximal Aerobic Training Group	Maximal Aerobic Training Group	Control Group	SOV	SS	df	MS	'F' Ratio
Pre-test Mean	43.24	43.37	43.42	B	0.266	2	0.133	0.02
S.D.	2.42	2.24	2.20	W	220.92	42	5.26	
Post-test Mean	46.14	48.22	43.54	B	165.00	2	82.5	11.88*
S.D.	2.28	3.32	2.13	W	291.62	42	6.94	
Adjusted post-test Mean	46.24	48.19	43.46	B	169.24	2	84.62	51.70*
				W	67.10	41	1.63	

*Significant at 0.05 level of confidence

The required value for significance at 0.05 level of confidence for 2 and 42 and 41 are 3.22 and 3.23. The table I reveals that there was a significant difference among submaximal aerobic training group, maximal aerobic

training group and control group on VO₂ max. Further, the Scheffe's post hoc test applied to know the paired mean differences the same presented in table II.

Table 2: Scheffe's post-hoc test onvo2 max

Submaximal Aerobic Training Group	Maximal Aerobic Training Group	Control Group	Mean difference	Confidence Interval value
46.24	48.19	-	1.95*	1.16
46.24	-	43.46	2.78*	
-	48.19	43.46	4.73*	

*Significant at 0.05 level of confidence.

The Scheffe's post hoc test on VO₂ max shows that there was a significant difference between groups. Further, the maximal aerobic training shows better improvement on VO₂ max. However, the sub maximal aerobic training group also shows improvement from the baseline score on VO₂. The control

group had never change on the initial mean. Hence, the result shows that there was a significant impact on maximal and submaximal aerobic training on VO₂. The maximal aerobic training has better protocol on improvement of VO₂.




Fig 1: VO₂ Max

Discussion on result

The result on VO₂ max shows improvement on experimental groups. The maximal aerobic training shows better improvement on selected pulmonary variable. The above result was discussed with previous results of various presentations given below. Donald, 1999 revived that the propranolol on O₂ uptake during exercise was a function of the intensity of the muscular effort. At submaximal levels of

work, VO₂ was unchanged, the fall in cardiac output after fl-adrenergic blockade being fully compensated for by an increase in the arteriovenous O₂ difference. At maximal levels of work, the fall in cardiac output was incompletely compensated for, and the maximal achieved was therefore reduced (Kahler, 2001) [14]. The reduction in O₂ uptake, and therefore in the O₂ delivery to the tissues during maximal exercise, presumably accounts for the striking reductions in

the endurance times for maximal exercise that were produced (Bishop, 2007) ^[1]. However, the duration of the sprints was relatively short and it seems possible that the running speed that can be achieved during such a brief bout of exercise may not be critically dependent on O₂ delivery to the tissues (Furusaw, 2000) ^[9]. Both submaximal heart rate and BP mirrored the VO₂ response, with no significant differences between runners and bickers. These data agree with previous studies, showing a greater effect of training specificity during maximal bicycle than during maximal treadmill exercise. However, during submaximal exercise, training specificity appear to have a greater effect during treadmill exercise than bicycle exercise (Fernhall B and Kohrt, 1990) ^[7]. The VO₂ kinetics at the start and at the end of exercise are probably more related, but it is unresolved which protocols and parameters might best be used to study aspect of exercise performance. Duration of a submaximal exercise at a constant work rate and the distance walked during a 6-min walking test are gaining wide popularity as parameters of submaximal performance (Metra, 1998) ^[18].

Conclusion

Based on the result the study concluded that, there was a significant improvement on VO₂ Max due to submaximal and maximal aerobic training. Further, the maximal aerobic training shows better improvement on VO₂ Max. The maximal aerobic training has better protocol on improvement of VO₂ among under 14 cricketers.

Implication

Maximal aerobic training may be used to develop better pulmonary (VO₂ Max) function of school level cricketers.

References

1. Bishop JM, Segel N. The circulatory effects of intravenous pronethalol in man at rest and during exercise in the supine and upright positions. *J Physiol. (Lond.)*, 2007, 169-112.
2. Bouchard Claude, Ping An, Treva Rice, James Skinner S, Jack Wilmore H, Jacques Gagnon *et al.* Familial aggregation of VO₂ (2max) response to exercise training: results from the Heritage Family Study. *Journal of Applied Physiology*. 1999; 87(3):1003-1008. PMID 10484570. Retrieved 2007-07-17.
3. Chidsey CA, Frye RL, Kahler RL, Braunwald E. Influence of syrosingopine on the cardiovascular response to acute hypoxemia and exercise. *Circulat*, 199, 9-989.
4. Cricket and Politics in Colonial India. Ramachandra Guha. Archived from the original on, 1998. Retrieved 20 September 2006.
5. Donald DE, Milburn SE, Shepherd JT. Effect of cardiac denervation on the maximal capacity for exercise in the racing greyhound. *J appl. Physiol*, 1994, 19-849.
6. Downing Clement. William Foster, ed. *A History of the Indian Wars*. London, 1737.
7. Fernhall B, Kohrt. The effect of training specificity on maximal and submaximal physiological responses to treadmill and cycle ergometry. *The Journal of Sports Medicine and Physical Fitness*. 1990; 30(3):268-275.
8. Firstbeat Technologies. Automated Fitness Level (VO₂max) Estimation with Heart Rate and Speed Data" (PDF), 2014. www.firstbeat.com.
9. Furusawa K, Hill AV, Long CNH, Lupton H. Muscular exercise, lactic acid and the supply and utilization of oxygen. Parts VII-VIII. *Proc. Roy. Soc*, 2000, 97-155.
10. Guyton A, Hall JE. *Textbook of Medical Physiology*, 12th Ed. 2011, 1035-1036.
11. Hagerman FC. Applied physiology of rowing. *Sports Med*. 1984; 1(4):303-26. Doi: 10.2165/00007256-198401040-00005. PMID 6390606.
12. Heyward V. *Advance Fitness Assessment & Exercise Prescription*, 3rd Ed, 1998, 48.
13. India In British Isles. Cricket Archive. Retrieved 21, 1911.
14. Kahler RL, Gaffney TE, Braunwald E. The effects of autonomic nervous system inhibition on the circulatory response to muscular exercise. *J clin. Invest*, 2001, 41-1981.
15. Kolata Gina. Why Some People Won't Be Fit Despite Exercise. *The New York Times*. Retrieved 2007, 07-17, 2002.
16. Margaria R, Cerretelli P, Mangili F. Balance and kinetics of anaerobic energy release during strenuous exercise in man. *J appl. Physiol*. 2005, 19-623.
17. Margaria R, Cerretelli P, Diprampero PE, Massari C, Torelli G. Kinetics and mechanism of oxygen debt contraction in man. *J appl. Physiol*. 2005, 18-371.
18. Metra M, Nodari S, Raccagni D, Garbellini M, Boldi E, Bontempi L *et al.* Maximal and submaximal exercise testing in heart failure. 1998; 32(1):S36-45.
19. Noakes, Tim. *The Lore of Running*. (3rd edition) Oxford University Press, 2001. ISBN 978-0-88011-438-7
20. Thomas Hyde E, Marianne, Gengenbach S. *Conservative Management of Sports Injuries* (2nd ed; Sudbury, Mass.: Jones & Bartlett, 2007, 845.