

ISSN: 2456-0057  
 IJPNPE 2019; 4(2): 542-545  
 © 2019 IJPNPE  
 www.journalofsports.com  
 Received: 07-05-2019  
 Accepted: 10-06-2019

**U Prabhakari**  
 Ph.D., Scholar Part Time,  
 Annamalai University,  
 Chidambaram, Tamil Nadu,  
 India

**Dr. V Gopinath**  
 Registrar, Tamil Nadu Physical  
 Education and Sports  
 University, Chennai, Tamil  
 Nadu, India

## Effect of asanas and pranayama on selected biochemical variables among polycystic ovary syndrome

U Prabhakari and Dr. V Gopinath

### Abstract

The purpose of the study was to find out the effect of asanas and pranayama on selected biochemical variables among polycystic ovary syndrome. To achieve this purpose 45 polycystic ovary syndrome women were randomly selected from Trichy district, Tamil Nadu and their age ranged between 25-35 years. They were divided in to three equal groups, namely Experimental Group I (Asanas), Experimental Group II (Pranayama) and Control Group. The training period was 12 weeks and 6 days per week. No treatment was given to control group and kept separate. High density lipoproteins and low density lipoproteins were selected as dependent variables. The pre and post test was conducted before and after the treatment respectively. The ANCOVA was used to find out the significant difference between the groups. If the result was significant Scheffe's post hoc test was used to find out the paired mean differences. The 0.05 level of confidence was fixed to find out the significant level. It was concluded that effect of asana and pranayama improves the biochemical variables such as triglycerides and high density lipoproteins.

**Keywords:** Asanas, pranayama, HDL and LDL

### Introduction

Polycystic ovary syndrome (PCOS) is a complex heterogeneous endocrine disorder. It is a common disorder affecting 4-12% of women of reproductive age (Sheehan, 2004; Azziz, *et al.* 2004) [14, 3]. PCOS was first described in the United States in 1935 (Stein and Leventhal, 1935) [15]. PCOS is characterized by chronic anovulation and hyper androgenism in the absence of underlying adrenal or pituitary disease. Women with PCOS may complain about variable clinical manifestations including oligomenorrhea, hirsutism, acne, and infertility (Ehrmann, 2005) [8]. Approximately 75% of these women suffer from infertility due to anovulation. Therefore, it is the most common cause of an ovulatory infertility (Adams, Polson and Franks, 1986; Hull, 1987) [1, 11]. PCOS is also reported to be associated with obesity, insulin resistance and type II diabetes, dyslipidemia, hypertension, cardiovascular disease and endometrial carcinoma (Carmina and Lobo, 1999) [4]. Approximately 50-60% of women with the syndrome are overweight or obese compared to 30% of women in the general population (McKittrick; Flegal, *et al.* 2002) [12, 9]. Treatment of PCOS must focus both on normalizing short-term signs of hyper androgenism and anovulation and on reducing metabolic complications. This can be achieved through pharmacological intervention or preferably lifestyle modification (Norman, *et al.* 2002) [13]. The most preferred and effective method of treatment of PCOS is lifestyle modification. Weight loss is an important treatment strategy. Weight loss improves practically every parameter of PCOS. In obese, an ovulatory PCOS women, weight loss restores ovulation and pregnancy rates, decreases insulin levels, diminishes acanthosis nigricans, lowers testosterone levels while raising sex hormone binding globulin (SHBG) levels, and improves psychological considerations (Clark, *et al.* 1995; Apter, 1998) [5, 2]. Despite the potential beneficial effects of exercise in PCOS, there are no systematic reviews that evaluate the independent effects of exercise on cardiovascular and reproductive which are supposed to play a significant role in the future performance of a players such as Despite the potential beneficial effects of exercise in PCOS there are no systematic reviews that evaluate the independent effects of exercise on cardiovascular and reproductive outcomes. Two recent non-systematic exercise-specific reviews highlight the gaps in knowledge around exercise type

**Corresponding Author:**  
**U Prabhakari**  
 Ph.D., Scholar Part Time,  
 Annamalai University,  
 Chidambaram, Tamil Nadu,  
 India

and intensity required to improve outcomes in PCOS (Hoeger, 2008; Thomson *et al.* 2010) [10, 16]. Consideration of the frequency and duration of exercise, as well as an assessment of sustainability, is also needed to determine the role of exercise in PCOS. To our knowledge is to evaluate the prevalence of PCOS in women. Our main objective was to estimate the bio chemical variables of PCOS in physically active women and to determine the effect of asanas and pranayama.

**Statement of the problem**

To find out the effect of asanas and pranayama on selected biochemical variables among polycystic ovary syndrome.

**Methods and Materials**

A total of 45 women with polycystic ovary syndrome were selected at random from Trichy District, Tamil Nadu and their age ranged between 25-35 years and they were divided in to

three groups consisting of fifteen subjects namely Experimental Group I, Experimental Group II and Control Group. Experimental group I underwent asanas, experimental group II underwent pranayama for a period of 12 weeks and 6 days per week. No treatment was given to control group and kept separate. High density lipoproteins and low density lipoproteins were selected as the dependent variables. The study was formulated true random group design, consisting of a pre-test and post-test. The data were collected before and after the treatment respectively. The pre-test and post-test scores were collected and the obtained data were subjected to statistical treatment using analysis of covariance (ANCOVA). If the adjusted post-test result was significant, the Scheffe's post hoc test was used to determine the significance of the paired mean differences. In all cases the level of significance at 0.05 level of confidence was fixed as appropriate.

**Results and discussion on high density lipoproteins**

**Table 1:** Computation of Analysis of Covariance on HDL (Scores in mg/dl)

Means	Exp. Group I	Exp. Group II	Control Group	SV	SS	df	MS	Obtained 'F'
Pre test	35.60	36.60	36.93	B	14.44	2	7.22	0.55
				W	548.13	42	13.05	
Post test	41.67	39.47	36.87	B	173.20	2	86.60	8.21*
				W	442.80	42	10.54	
Adjusted post test	42.16	39.32	36.51	B	233.99	2	116.99	21.96*
				W	218.46	41	5.33	

\*Significant at 0.05 level. Table value required for significance at 0.05 level for df 2 and 42 is 3.22 and df 2 and 41 is 3.23.


**Table 2:** Computation of Scheffe's Post Hoc Test of HDL (Scores in mg/dl)

Exp. Group I	Exp. Group II	Control Group	MD	CI
42.16	39.32	-	2.84*	2.14
42.16	-	36.51	5.65*	2.14
-	39.32	36.51	2.81*	2.14

obtained F value 0.55, 8.21 and 21.96 respectively. This result clearly indicated that the effect of asanas and pranayama significantly increased the high density lipoprotein level among women with polycystic ovary syndrome in post and adjusted posttest. Scheffe's Post Hoc test was given in the table II indicated that the ordered adjusted final mean difference between groups.

**Discussion on high density lipoprotein**

The table I showed pre, post and adjusted posttest mean



**Fig 1:** Bar Diagram Showing the Pre and Post Test Means of High Density Lipoprotein (Scores in mg/dl)

## Results and Discussion on low density lipoproteins

**Table 3:** Computation of Analysis of Covariance on LDL (Scores in mg/dl)

Means	Exp. Group I	Exp. Group II	Control Group	SV	SS	df	MS	Obtained 'F'
Pre test	82.73	77.47	80.67	B	211.24	2	105.62	1.81
				W	2454	42	58.43	
Post test	74.87	72.87	80.87	B	520	2	260	5.54*
				W	1971.20	42	46.93	
Adjusted post test	72.94	75.09	80.57	B	463.152	2	231.58	20.93*
				W	453.730	41	11.07	

\*Significant at 0.05 level. Table value required for significance at 0.05 level for *df* 2 and 42 is 3.22 and *df* 2 and 41 is 3.23.


**Table 4:** Computation of Schaffer's Post Hoc Test of LDL (Scores in mg/dl)

Control Group	Exp Group II	Exp Group I	MD	CI
80.57	75.09	-	5.48*	3.09
80.57	-	72.94	7.63*	3.09
-	75.09	72.94	2.15	3.09

\*Significant at 0.05 level of confidence

### Discussion on low density lipoproteins

The table III showed pre, post and adjusted posttest mean obtained F value 0.55, 8.21 and 21.96 respectively. This results clearly indicated that the effect of asanas and pranayama significantly decreased the low density lipoproteins level among women with polycystic ovary syndrome in post and adjusted posttest. Scheffe's Post Hoc test was given in the table IV indicated that the ordered adjusted final mean difference between groups.



**Fig 2:** Bar Diagram Showing the Pre and Post Test Means of Low Density Lipoprotein (Scores in mg/dl)

### Conclusion

PCOS is a prevalent disorder which is underpinned by weight independent IR and associated with reproductive and cardiovascular complications in women. Lifestyle modification, including exercise, is the first-line therapeutic approach in improving health outcomes in PCOS. Despite this, the present review identified exercise intervention in PCOS. Results from this review suggest that improvements in dependent variables such as high density lipoproteins and low density lipoproteins on length of asanas and pranayama intervention as significant benefits were observed in more sustainable of shorter duration. Regular, asanas and pranayama improves reproductive outcomes including ovulation and menstrual cycle regulation in addition to reducing weight women with PCOS. Future studies would benefit by using gold standard, comprehensive end points to provide mechanistic insights into PCOS particularly central

feature of this endocrinopathy. Based on the results from this study, women with PCOS should be advised to engage in at least 90 min of asanas and pranayama six days per week to achieve improved reproductive and cardio metabolic outcomes.

### Reference

- Adams J, Polson DW, Franks S. Prevalence of polycystic ovaries in women with anovulation and idiopathic hirsutism. *BMJ*. 1986; 293:355-359.
- Apter D. How possible is the prevention of polycystic ovary syndrome development in adolescent patients with early onset of hyper androgenism. *J Endocrinol Invest*. 1998; 21:613-617.
- Azziz R, Woods KS, Reyna R *et al*. The prevalence and features of the polycystic ovary syndrome in an unselected population. *J Clin Endocrinol Metabolism*. 2004; 89:2745-2749.
- Carmina, E, Lobo RA. Polycystic ovary syndrome (PCOS): arguably the most common endocrinopathy is associated with significant morbidity in women. *J Clin Endocrinol Metab*. 1999; 84:1897-1899.
- Clark AM, Ledger W, Galletly C *et al*. Weight loss results in significant improvement in pregnancy and ovulation rates in an ovulatory obese women. *Hum Reprod*. 1995; 10:2705-2712.
- Dahlgren E, Janson PO, Johansson S *et al*. Polycystic ovary syndrome and risk for myocardial infarction. Evaluated from a risk factor model based on a prospective population study of women. *Acta Obstet Gynecol Scandinav*. 1992; 71:599-604.
- Dunaif A, Segal KR, Futterweit W *et al*. Profound peripheral insulin resistance, independent of obesity, in polycystic ovary syndrome. *Diabetes*. 1989; 38:1165-1174.
- Ehrmann DA. Polycystic Ovary Syndrome. *N Engl J Med*. 2005; 352:1223-1236.
- Flegal KM, Carroll MD, Ogden CL *et al*. Prevalence and trends in obesity among US adults. 1999-2000; 2002:1772-1773.
- Hoeger KM. Exercise therapy in polycystic ovary syndrome. *Semin Reprod Med*. 2008; 26:93-100.
- Hull MGR. Epidemiology of infertility and polycystic ovarian disease: endocrinological and demographic studies. *Gynecol Endocrinol*. 1987; 1:235-245.
- McKittrick M. Diet and polycystic ovary syndrome. *Nutria Today*. 2002; 37:63-69.
- Norman R, Davies M, Lord J *et al*. The role of lifestyle modification in polycystic ovary syndrome. *TRENDS Endocrinol Metab*. 2002; 13:251-257.
- Sheehan MT. Polycystic ovarian syndrome: diagnosis and management. *Clin Med Res*, 2004; 2:13-27.

15. Stein IF, Leventhal NL. Amenorrhea associated with bilateral polycystic ovaries. *Am J Obstet Gynecol.* 1935; 29:181-191.
16. Thomson RL, Buckley JD, Brink worth GD. Exercise for the treatment and management of overweight women with polycystic ovary syndrome: a review of the literature. *Obese Rev*, 2010.