



ISSN: 2456-0057

IJPNPE 2023; 8(1): 303-304

© 2023 IJPNPE

www.journalofsports.com

Received: 04-02-2023

Accepted: 16-03-2023

Dr. G Jayapal

Assistant Professor, Meenatchi
Physical Education College,
Thathanur, Tamil Nadu, India

Dr. A Nallamuthu

Assistant Professor,
Dhanalakshmi Srinivasan
College of Physical Education
Perambalur, Tamil Nadu, India

Impact of anaerobic training on strength endurance among college men students

Dr. G Jayapal and Dr. A Nallamuthu

Abstract

The purpose of present study was to find out the impact of anaerobic training on strength endurance among college men students. To achieve this purpose, thirty men students chosen from Meenatchi Physical Education College, Thathanur, Tamil Nadu, India were selected as subjects. The selected thirty subjects were randomly divided into two groups of fifteen each, out of which group - I (n = 15) underwent anaerobic training (short bursts of high intensity movement) for three days (alternative days) per week, group - II (n = 15) remained as control. Prior to and after the training period the subjects were tested for, strength endurance. strength endurance was assessed by bent knee sit-ups test. The statistical tool were used for the present study is 't' ratio and Analysis of covariance (ANCOVA). The result of the study was a significant altered on strength endurance after twelve weeks of anaerobic training. However the different was favour of experimental group.

Keywords: anaerobic training, strength endurance

Introduction

Physical training aims at improving the performance of sports persons. Several factors influence ties sports performance. The performance of sports primarily depends on his performance capacity, such as speed, strength and endurance. All these factors therefore are the principal aims of physical training.

Anaerobic means 'without oxygen'. During anaerobic work, involving maximum effort, the body is working so hard that the demands for oxygen exceed the rate of supply and the muscles have to rely on the stored recoveries of fuel. In this case waste product accumulate, the chief one being lactic acid. The muscles, being starved of oxygen, take the bodies into a state known as oxygen debt. The body's stored fuel soon runs out and activity ceases with pain.

Many workout routines utilize periods of intense activity as a part of their regimen. Increasing anaerobic capacity has been shown to have a number of health benefits, including better athletic performance and increased metabolism.

Muscular strength is the amount of force you can put out or the amount of weight you can lift. Muscular endurance is how many times you can move that weight without getting exhausted (very tired).

Statement of the problem

The purpose of present study was to find out the effect of anaerobic training on strength endurance among college men students.

Methodology

The purpose of present study was to find out the impact of anaerobic training on strength endurance among college men students. To achieve this purpose, thirty men students chosen from Meenatchi Physical Education College, Thathanur, Tamil Nadu, India were selected as subjects.

Corresponding Author:

Dr. G Jayapal

Assistant Professor, Meenatchi
Physical Education College,
Thathanur, Tamil Nadu, India

The selected thirty subjects were randomly divided into two groups of fifteen each, out of which group - I (n = 15) underwent anaerobic training (short bursts of high intensity movement) for three days (alternative days) per week, group - II (n = 15) remained as control. Prior to and after the training period the subjects were tested for, strength endurance. strength endurance was assessed by bent-knee sit-ups test.

Analysis of data

Table-I presents pre and post-test means, standard deviations and dependent t-test values on strength endurance of anaerobic training and control group.

Table 1: Summary of means, standard deviation and dependent t-test values on strength endurance of anaerobic training and control group

Tests	Anaerobic Training Group		Control Group	
	Mean	SD	Mean	SD
Pre test	33.25	2.01	33.26	2.06
Post test	37.42	2.11	33.21	0.03
T-Test	5.32*		0.52	

*Significant at .05 level. The table value required at .05 level with df 14 is 2.14.

From the table-I shows that the obtained dependent t-test values between pre-test and post-test means of anaerobic training and control group are 5.32 and 0.51 respectively. The table value required for significant difference with df 14 at .05 level is 2.14. Since, the obtained t-test value of anaerobic training group is greater than the table value, it is understood that anaerobic training programme had significantly improved the performance of strength endurance and the control group has not improved as the obtained t-test value is lesser than the table value because they were not subjected to any specific training. The analysis of covariance on strength endurance of anaerobic training and control group have been analysed and presented in table - 2

Table 2: Analysis of covariance on strength endurance of anaerobic training and control group

Adjusted post-test mean		Source of variance	Sum of squares	Df	Mean square	F-ratio
Anaerobic Training Group	Control Group	Between	0.897	1	0.695	28.65*
36.89	33.21	Within	0.323	27	0.22	

*Significant at .05 level of confidence. The table value required at 0.05 level with df 1 & 27 is 4.21.

Table- 2 shows that the adjusted post-test means of anaerobic training and control groups are 36.89 and 33.21 respectively. The obtained f-ratio value is 28.65 which is higher than the table value 4.21 with df 1 and 27 required for significance at .05 level.

It was concluded that anaerobic training is better than control groups in improving strength endurance.

Conclusions

The anaerobic training group has remarkably increase strength endurance when compared with the control group. In addition, the results of the tests shows that there was significant difference established between experimental and control groups.

References

1. Petr  H, Wernst l F, Mattsson CM. Effects of Flywheel Training on Strength-Related Variables: a Meta-analysis.

In Sports Medicine – Open; c2018, 4(1). <https://doi.org/10.1186/s40798-018-0169-5>

- Raja DSC. Influence of resistance and endurance training and combination of resistance and endurance training on back strength flexibility and cardio-respiratory endurance. International Journal of Physical Education, Sports and Health; c2021, 8(3). <https://doi.org/10.22271/kheljournal.2021.v8.i3e.2099>
- Sch rer C, Tacchelli L, G pfert B, Gross M, L thy F, Taube W, *et al.* Specific eccentric–isokinetic cluster training improves static strength elements on rings for elite gymnasts. International Journal of Environmental Research and Public Health; c2019, 16(22). <https://doi.org/10.3390/ijerph16224571>
- Schwiete C, Franz A, Roth C, Behringer M. Effects of Resting vs. Continuous Blood-Flow Restriction-Training on Strength, Fatigue Resistance, Muscle Thickness, and Perceived Discomfort. Frontiers in Physiology; c2021, 12. <https://doi.org/10.3389/fphys.2021.663665>